

Treatment of Misophonia

M. Zachary Rosenthal, PhD^{a,b,*}, Yanyan Shan, MA^{a,b}, Jacqueline Trumbull, MA^{a,b}

^aDepartment of Psychiatry & Behavioral Sciences, Duke University, DUMC, 6400 Pratt Street, North Pavilion, 6th Floor, Room 6043, Durham, NC 27705, USA; ^bDepartment of Psychology & Neuroscience, Duke University, Durham, NC, USA

KEYWORDS

- Misophonia
- Treatment
- Psychotherapy
- Unified protocol
- Process-based therapy
- Multi-disciplinary treatment

KEY POINTS

- Introduces misophonia and reviews published psychotherapy treatment studies.
- Outlines a multi-disciplinary strategy for treatment.
- Describes the application of two transdiagnostic psychotherapies with emerging evidence in misophonia (Unified Protocol and Process-Based Therapy).
- Suggests an agenda for future research and treatment development.

Misophonia was recently defined as a disorder characterized by intolerance of specific auditory stimuli and associated cues causing significant psychological distress and interference in social, occupational, or academic functioning [1]. Aversive stimuli in this context may be labeled as “triggers” that are commonly repetitive sounds produced by others. Primary triggering stimuli in misophonia most often (but not always) originate from facial (eg, nose noises, throat-clearing) or oral (eg, lip-smacking, chewing, drinking) sources. However, there are individual differences in the types of cues and contexts (eg, the same sound may have different effects when produced by specific people) associated with misophonia symptoms [2,3].

Individuals with impairing levels of misophonia symptoms may have strong multi-modal emotional responses to contexts in which triggering cues are anticipated or encountered. For example, common responses include central (eg, insula) [4] and peripheral nervous system activation (eg, increased heart rate and skin conductance) [5], negative affect (eg, irritation, anger, anxiety, disgust) [6,7], and behavioral patterns that can

be conceptualized as consistent with freeze (eg, hypervigilance), flight (eg, escape or avoidance behavior), and fight behaviors (eg, interpersonally aggressive verbal behavior) [8]. These responses are highly distressing and different than how most others might react [1].

Jastreboff and Jastreboff are credited for coining the term misophonia over 20 years ago [9]. The term translates to hatred or dislike (miso) of sound (phonia). However, this literal translation is misleading, as the condition is neither uniquely associated with the affective experience of hate or dislike, nor are auditory cues the only stimuli that can function as triggers. The first pilot studies directly examining misophonia were published only 9 years ago [5,10]. Since then, over 60 published empirical studies have investigated misophonia, with roughly 25% of these published over the last year in the first special section in a peer-reviewed scientific journal dedicated to misophonia (*Frontiers in Neuroscience*).

Recent reviews suggest that misophonia may be associated with a wide variety of mental health problems [1–3]. The majority of studies have used self-report

*Corresponding author, E-mail address: mark.rosenthal@duke.edu

methodologies and have found that misophonia symptom severity is positively correlated transdiagnostically with, for example, neuroticism, anxiety symptoms, depressive symptoms, difficulties with emotion regulation, affective instability, anxiety sensitivity, some obsessive-compulsive disorder (OCD) symptoms, perfectionism, and somatic pain [11–19]. Additionally, adults with misophonia may be significantly more likely than those without misophonia to self-report a lifetime history of attention-deficit/hyperactivity disorder (ADHD), OCD, bipolar disorder, substance use disorder, post-traumatic stress disorder, and conversion disorder [17,20].

Two recent studies have used structured psychiatric diagnostic interviews as a more rigorous approach to the assessment of misophonia and associated mental health problems. In one study, 575 adults presenting for treatment at a clinic in Amsterdam were interviewed using the Mini International Neuropsychiatric Interview [21], a structured interview assessing 15 current psychiatric problems (Jager and colleagues, 2020a) [14]. Results indicated that most (72%) participants did not meet the full criteria for a current psychiatric disorder. The most common current disorders were mood disorders (10.1%), anxiety disorders (9%), ADHD (5.4%), and personality disorders (5%).

Rosenthal and colleagues [8] conducted the first study to characterize diagnostic and statistical manual of mental disorders (DSM-5) disorders using structured diagnostic interviews in a nationally recruited community sample of 207 adults with high misophonia symptoms. Results indicated that anxiety disorders were the most common type of current mental health problem (56.9%). Additionally, high rates of lifetime history of psychiatric disorders were observed, including diagnoses of any anxiety (73%), mood (61%), obsessive-compulsive (27%), substance use (26%), trauma-related (24%), eating (18%), or personality disorders (13%).

The findings from Jager and colleagues [14] suggest that most adults seeking outpatient treatment of misophonia in Amsterdam may not have a current psychiatric disorder. In contrast, Rosenthal and colleagues [8] indicate that adults across the United States with high misophonia symptoms may be most likely to meet the full criteria for a current anxiety disorder or lifetime history of any anxiety or mood disorders. A consistent interpretation of findings across all studies is that higher misophonia symptoms do not appear to be uniquely related to any specific psychiatric disorder. Accordingly, it is premature to conclude that any specific treatment protocol for any specific psychiatric disorder is best for the treatment of misophonia.

TREATMENT OF MISOPHONIA

Multi-Disciplinary Model: Audiology

There is no gold standard evidence-based treatment of misophonia, and it is not a disorder within the purview of any specific clinical discipline. Because it appears to be a problem at the intersection of various clinical fields, a multi-disciplinary treatment model may be valuable. For example, audiologists may be important in the assessment and treatment of misophonia as it is defined as a sound intolerance condition. Because misophonia may need to be differentiated from hyperacusis, audiologists may be helpful in assessment and treatment planning. For example, audiologists use evaluative measures to assess sensitivity thresholds and can discern if one is hypersensitive to auditory stimuli.

With regard to treatment, audiologists can work with patients to determine the pros and cons of using sound-based therapies. These approaches use patient-controlled devices unobtrusively placed in the ear canal to diminish (eg, noise cancellation), mask (eg, static brown or white noise), and change responses to auditory input. Although these interventions have not been tested empirically using randomized trials for the treatment of misophonia, promising support has come from clinical observations reported by Jastreboff and Jastreboff, who pioneered the adaptation of Tinnitus Retraining Therapy for patients with misophonia and report high rates of success in uncontrolled trials [22]. This approach uses sound therapy and behavioral training to change response patterns to triggering contexts and cues, and was reported by the developers to be highly effective in a clinical setting.

Multi-Disciplinary Model: Occupational Therapy

Because misophonia may occur in the context of multi-sensory over-responsivity, it may be valuable to include occupational therapists in a multi-disciplinary assessment and treatment approach. Occupational therapists emphasize improvement of functioning across key domains of life. Because occupational therapists are the experts in sensory processing within health care, they may offer helpful coping strategies to manage misophonia effectively. Occupational therapy includes treatments designed to help improve central sensory processing through interventions designed to enhance sensory integration functioning. Treatments may identify and intervene upon environmental barriers to adaptive sensory processing or train new adaptive reactions to emotionally evocative sensory cues. Like approaches used by audiologists, there are no randomized

controlled trials evaluating the efficacy of occupational therapy interventions with misophonia. However, these interventions have a long history of being used to improve sensory processing, and, as such, may be worth considering for those with misophonia.

Multi-Disciplinary Model: Mental Health

Some early clinical observations [22] and the large study by Jager and colleagues [14] indicate that adults with misophonia may present for treatment without co-occurring psychiatric disorders. On the other hand, there is a growing body of research and clinical observations pointing to misophonia occurring in the context of varied and debilitating psychiatric disorders [8] and an associated need for mental health professionals to be involved in the treatment of this disorder. Accordingly, evaluation and treatment recommendations from mental health providers are indicated in a multi-disciplinary model of care for misophonia.

Treatment Studies

Most publications describing possible psychotherapies for misophonia have been small case studies (Table 1) [23–35]. These treatments were mostly conducted using branded (eg, acceptance and commitment therapy; ACT) and non-branded interventions from the family of cognitive behavioral therapies (CBTs), and the authors reported descriptions of successful treatment in one or several individuals.

Examples of non-branded interventions include cognitive restructuring [23,24,26,30–32,35], relaxation exercises [26,28,32,35], counter-conditioning [27,28], acceptance and distress tolerance strategies [25,26,29,30,33,35], exposure and response prevention [23–26,30–32,34], interpersonal communication skills [24,32,33], attentional control skills [24,30,34], and parent management training [26,32]. Case studies are useful beginning points for treatment development and provide a direction for clinicians in the absence of a clear gold standard of care. However, these case studies do not provide compelling evidence of efficacy for any particular intervention.

OPEN TRIALS

Three open trials have been conducted to treat misophonia [36–38]. Schröder and colleagues [38] conducted an uncontrolled trial involving 90 adults with misophonia, showing promise for a cognitive behavioral approach using brief group therapy. In this trial, 48% of participants improved on a clinician rating of outcome, and 30% reported a significant reduction in

symptoms on a self-report measure of misophonia. The treatment included four main components: (1) attentional shifting away from trigger stimuli, (2) counter-conditioning to disrupt classically conditioned associations between neutral/positive stimuli that have become paired with negative emotional experiences, (3) stimulus manipulation exercises (allowing the participant to manipulate trigger sounds), and (4) relaxation exercises. Because this intervention occurs in a group, validation and support from other sufferers may also be an important component of this treatment.

Frank and McKay reported that 18 participants received 12 sessions of exposure therapy before or after stress management training [36]. The exposure procedure used an inhibitory learning model, emphasizing altered expectations for the target sounds along with the deliberate practice of hearing sounds on an individual hierarchy. Rather than targeting habituation of psychological distress when exposed to triggers, the inhibitory learning approach to exposure enhances patient motivation to approach triggering cues and contexts by changing the features of these cues or their responses to them in an effort to change expectations, increase psychological flexibility, enhance valued actions, and increase the perception of control over reactions to misophonic cues. This study demonstrates that without habituation to certain sounds, inhibitory learning-based exposure procedures may be a promising way to enhance perceived control over emotional reactions to misophonia triggers.

Finally, Jager and colleagues [37] used an open trial to evaluate the efficacy of eye movement desensitization and reprocessing (EMDR) therapy in treating misophonia in eight participants. Misophonia-related emotionally disturbing memories were addressed with EMDR in an average of 2.6 sessions lasting 60 to 90 minutes. Participants reported a statistically significant 20% average reduction (with a large effect size estimate of $d = 1.14$) in symptoms of misophonia on a self-report measure of misophonia. No statistically significant changes were observed in psychological distress or impairment in functioning on secondary outcome measures. Due to the uncontrolled experimental design of these open trials, no definitive conclusions can be drawn from these studies about treatment efficacy, though these are valuable studies pointing to possible treatments to further evaluate.

RANDOMIZED CLINICAL TRIAL

Jager and colleagues [39] conducted the only randomized controlled trial for misophonia (Table 2). Using

TABLE 1
Case Studies of Misophonia Treatment

Studies	Treatment	Strategies								Age	Gender	Comorbidity	# of sessions	Misophonia Assessment		
		Cognitive Restructuring	Relaxation Exercises	Counter-Conditioning	Acceptance and Distress Tolerance	ERP	Interpersonal Skills	Attention Control	Parenting Training					Baseline	Post	Symptom Reduce
Bernstein [24], 2013	CBT	√				√	√		√	19	Female	None	6	NA	NA	NA
Dozier [27], 2015(a)	NRT			√						48	Female	NA	14	MAQ41	MAQ9	MAQ78%
Dozier [28], 2015(b)	NRT		√	√						21	Female	None	4	MAQ49	MAQ13	MAQ73%
McGuire [31], 2015	CBT	√				√				17	Female	None	10	MQ55 MSS12	MQ37 MSS7	MQ33% MSS42%
										11	Female	None	18	MQ31 MSS5	MQ25 MSS4	MQ19% MSS20%
Reid [34], 2016	CBT	√				√			√	14	Female	OCD, MDD, phobia, ADHD	14	AMISOS17	AMISOS7	AMISOS59%
Kamody [29], 2017	DBT				√					16	Female	social anxiety	71+35G	AMISO22 MAQ51	AMISO10 MAQ16	AMISO 55% MAQ 71%
Schneider [35], 2017	DBT and ACT		√		√					17	Male	None	10	AMISOS14	AMISOS8	AMISOS57%
Altýnöz 2018	CBT	√				√				18	Female	NA	6	MAS6 MPRS6 AMISOS11	MAS2 MPRS2 AMISOS4	MAS67% MPRS67% AMISOS64%
Muller [32], 2018	CBT	√	√			√	√		√	14	Female	None	24	NA	NA	NA
Dover [26], 2021	CBT	√	√		√	√			√	10	Female	OCSD	30	AMISOS10 MQ27	AMISOS3 MQ6	AMISO70% MQ78%
Lewin [30], 2021	UP	√			√	√			√	4 cases, NA			10	MAQ25 AMISOS13	MAQ13 AMISOS9	MAQ48% AMISOS31%
														MAQ18 AMISOS15	MAQ25 AMISOS12	MAQ+39% AMISOS20%
														MAQ12 AMISOS9	MAQ3 AMISOS6	MAQ75% AMISOS33%
														MAQ54 AMISOS17	MAQ13 AMISOS10	MAQ76% AMISOS42%
Cowan [25], 2022	EASE [1]				√	√				14	Female	NA	6	NA	NA	NA
Petersen [33], 2022	ACT				√	√				12	Female	NA	16	AMISOS10	AMISOS5	AMISOS50%

Abbreviations: ACT, acceptance and commitment therapy; ADHD, attention deficit hyperactivity disorder; AMISO, Amsterdam Misophonia Scale; CBT, cognitive behavioral therapy; DBT, dialectic behavioral therapy; EASE, experiential acceptance and stimulus engagement; ERP, exposure and reaction prevention; MAQ, misophonia assessment questionnaire; MAS, misophonia activation scale; MPRS, misophonia psychological response scale; MQ, misophonia questionnaire; MSS, misophonia severity scale; NA, not applicable; NRT, neural repatterning technique; OCD, obsessive-compulsive disorder; OCSD, obsessive-compulsive spectrum disorder; 71+35G, 7 individual sessions and 35 group sessions.

TABLE 2
Open Trials and Randomized Clinical Trial of Misophonia Treatment

Studies	Treatment	Cognitive Restructuring	Relaxation Exercises	Counter-Conditioning	Attention Control	ERP	Stimulus manipulation	Others	Participants	Age	Comorbidity	# of Sessions	Symptom Reduce
Open Trials													
Schröder [38], 2017	Group-CBT		√	√	√		√		90	36	NA	8	AMISOS-R 33%
Frank [36], 2019	Inhibitory Learning	√				√			18	35	56%	12	NA
Jager [37], 2021	EMDR							Desensitization and reprocessing	10	35	50% Axis I, 60% Axis II	2.6	AMISOS-R 20%
Randomized Clinical Trial													
Jager [39], 2020	Group-CBT	√	√	√	√		√	Psychoeducation for family and friends	27	31	NA	12	AMISOS-R 32%

Abbreviations: AMISOS-R, Amsterdam misophonia scale—revised; CBT, cognitive behavioral therapy; EMDR, eye movement desensitization and reprocessing; NA, not applicable.

a cross-over design, adults with misophonia ($N = 54$) were randomly assigned to 3 months of weekly CBT in a group therapy format or to a waitlist. Participants were assessed at baseline, 3 months (following CBT or waitlist), 6 months (after cross-over), and between months 15 and 18 (1-year follow-up). CBT groups included task concentration, arousal reduction, positive affect labeling, and stimulus manipulation. Across all participants, symptoms of misophonia reduced by 32% after 3 months of CBT (a large effect size estimate of $d = 1.97$). Clinical improvement was observed in 37% of the CBT group compared to 0% in the waiting list group. Those who responded to CBT at post-treatment demonstrated no significant changes in symptoms of misophonia 1 year later, suggesting the effects of CBT maintained over time.

This randomized trial is the clearest evidence supporting the use of any specific treatment of misophonia. However, it should be noted that the comparison condition was an unblinded wait list control without any active intervention components. Additional studies are needed comparing this protocol to credible control conditions featuring non-specific factors such as psychoeducation, therapist time and attention, validation, and/or support. Nonetheless, the findings of this trial and, collectively, the combined findings across the preliminary case studies and open trials largely indicate that components from CBTs (eg, cognitive restructuring, exposure and counter-conditioning, relaxation exercises, interpersonal communication, and acceptance-based skills) may be reasonable for clinicians to consider and in need of future study as treatments of misophonia in both youth and adults.

TRANSDIAGNOSTIC APPROACHES TO PSYCHOTHERAPY

Unified Protocol

As outlined, early studies suggest misophonia (a) is not best accounted for by any psychiatric disorder, (b) is associated with various transdiagnostic psychological problems, and (c) pilot studies developing promising treatments have used various interventions from the family of CBTs. Accordingly, when considering which treatment approaches to scientifically evaluate, it is unclear whether it is appropriate to use a specific branded psychotherapy protocol designed for a specific diagnosis. Although untested, transdiagnostic psychotherapies may be a flexible and pragmatic alternative. The Unified Protocol (UP) is a reasonable candidate for transdiagnostic psychotherapy to consider, as it has been developed

and evaluated for use with a wide range of “emotional disorders,” including anxiety and mood disorders [40].

The UP is a 16-week skills-based treatment that consists of five core modules: mindful emotional awareness, cognitive flexibility, identifying and changing emotional avoidance, increasing tolerance of emotion-related physical sensations, and emotional exposures [41]. Patients with misophonia may be characterized by problems with emotional reactivity, limited access to emotion regulation strategies, and intolerance of elevated physical sensations when exposed to trigger sounds [13,42]. The core modules in the UP target these processes. The transdiagnostic approach of the UP has demonstrated efficacy against single-disorder treatments in multiple studies [41,43]. This success may be due to the high rates of co-occurrence between emotional disorders, with anxiety and depressive disorders having lifetime rates of co-occurrence as high as 75% [43]. As examples of efficacy, the UP has demonstrated improvements in anxiety [41], OCDs [44], and dysregulated anger across a range of clinical presentations [45]. Patients with misophonia may have high rates of co-occurring mental health problems [8,14], suggesting that the UP may be a helpful approach to consider.

Although a recent case study examined UP as a possible treatment of adolescents with misophonia [46], controlled clinical trials are needed before conclusions can be made about the efficacy of UP for misophonia. Our own research team is currently conducting preliminary open trials and developing a manual using the UP to treat adults with misophonia. Of eight participants treated with the UP, none dropped out of the treatment, and all reported that they found the treatment helpful. Patients also reported that learning about their own unhelpful coping mechanisms was important, as was enhanced skill use for emotional responses to trigger sounds [47].

Process-Based Therapy (PBT)

Although manualized models of psychotherapy such as the UP may be helpful, another candidate approach that is transdiagnostic and highly flexible is the PBT framework [48]. In PBT, therapists leverage evidence-based therapeutic processes that are common across therapies (eg, strong therapeutic alliance, empathy, support, motivational enhancement) and specific procedures used across protocols for various diagnoses and problems (eg, the list of interventions in Table 1). PBT assessment features functional analyses to identify maladaptive and adaptive patterns and measurement-based care using both qualitative and psychometrically validated quantitative measures of functioning and change

processes. Additionally, targets for treatment are collaboratively selected in a sequence that is acceptable to patients and intended to impact other targets in a network of related change processes. This enables an emphasis on patient strengths and empowerment, with assumptions of non-linear change leading to iterative and flexible changes in targets and therapeutic procedures throughout treatment.

When using PBT for misophonia, a therapist and patient determine through functional analyses that there are problematic patterns before, during, or after being triggered across attentional (eg, hypervigilance), cognitive (eg, internal, stable, and global attributions), physiological (eg, sympathetic arousal), social (eg, verbal confrontations), or other behavioral (eg, avoidance or escape behavior) levels of functioning. After discerning specific personally-relevant patterns, problematic patterns are collaboratively prioritized for targeted change. Next, therapeutic procedures known empirically to impact targeted patterns are offered by the therapist, and the patient chooses the one they are most willing and able to do (eg, to reduce physiological arousal when triggered, the patient could choose any protocol known to reduce sympathetic arousal). The intervention is administered, the patient applies the intervention in their daily life, and measures are taken to determine the effects of the intervention. When progress is made on a prioritized target, the patient selects the next most prioritized target to address, and the process repeats itself until patients are satisfied that they have met treatment goals. PBT is an approach that our group is currently testing for misophonia, with a manual under development iteratively based on patient feedback. As such, it is unknown if this approach is efficacious for misophonia.

RESEARCH AGENDA FOR TREATMENT DEVELOPMENT

Mechanistic Translational Studies

Studies identifying underlying neurobehavioral mechanisms of misophonia are needed to develop optimal interventions that target precise biological, social, or behavioral change processes. Candidate targets include difficulties with attention (eg, hypervigilance toward possible misophonic cues), cognition (eg, attributional styles, hopelessness), behavior (eg, avoidance, escape), social (eg, indirect aggression), and emotional (eg, sensitivity, reactivity) processes. As researchers seek to discover underlying biopsychosocial change processes in misophonia, it will be important to do so using multi-method studies with objective laboratory-based

measures. Processes that are unique to misophonia and those shared with other sound intolerance or mental health conditions need to be elucidated empirically using thoughtful experimental and statistical approaches.

Global Research with Diverse Participants

To date, most studies investigating misophonia have included disproportionately White, educated women as participants. It may be that these demographic factors align with access to knowledge about misophonia or motivation to participate in research. However, some studies using sampling methodologies with more representative approaches have not reported gender differences in misophonia symptoms [49]. Similarly, no studies have explored ethnicity, race, or multicultural considerations related to misophonia. To understand the nature and features of misophonia for all people, it is essential for researchers to use sampling methods that include diverse participants.

Multi-Disciplinary Treatment Models

Although a multi-disciplinary model of evaluations and treatment is recommended as a general strategy, there are no studies empirically testing this approach. To determine for whom, how, and why this model may be helpful, multi-disciplinary research is needed. This could include direct testing of such an approach compared to usual treatment or discipline-specific interventions. This also could include using adaptive designs that begin with fewer resources and, for those who are non-responsive, randomizing participants to higher resource treatment approaches.

Psychoeducation and Support

It is uncommon for clinical providers or the lay public to have extensive knowledge about misophonia. This overall lack of public awareness about misophonia frequently translates into patients and loved ones needing foundational psychoeducation and support in how to make sense of and manage misophonia. Particularly for caregivers and those with somewhat less impairing misophonia presentations, it may be that psychoeducation and support can serve as an important, low-cost, and scalable component of an overall treatment approach. Studies are needed to examine whether, for whom, and how such approaches can be helpful.

Digital Health

Another recommended approach for treatment development is the use of digital health-based models of care. This could include, for example, interventions using mobile phones with misophonia support apps to screen,

educate, support, and provide specific real-time interventions targeting underlying mechanistic change processes (eg, attentional hypervigilance, emotional reactivity, indirect interpersonal aggression, approach or avoidance behavior, cognitive reframing, or defusion). In addition, digital health approaches that train novel coping skills in virtual environments or with augmented reality could directly target underlying mechanistic targets of change and, in some instances, provide real-time feedback and tailored interventions based on user input and machine learning. These platforms can provide immersive and engaging experiences that may be more desirable for some than conventional treatment approaches such as psychotropic medication or psychotherapy.

CLINICS CARE POINTS

- Misophonia symptoms can occur with or without psychiatric disorders or other health problems. Start treatment with a multi-disciplinary strategy of evaluations and treatment recommendations across audiology, occupational therapy, and mental health providers.
- There are no cures, evidence-based medications, or proven treatments for misophonia. Aim to enhance functioning using interventions known to impact primary processes underlying problematic patterns (physiological arousal, attention, cognition, behavior, communication).
- Given the absence of a proven mental health treatment of misophonia, transdiagnostic evidence-based therapies (eg, UP) and flexible patient-centered frameworks leveraging evidence-based change process may be useful to tailor interventions to the individual (eg, PBT), although require further evaluation.
- Habituation-based exposure therapy is not indicated, but inhibitory learning models of exposure therapy may be helpful as part of a broader treatment approach.

FUNDING

Funding for this work was provided by anonymous supporters of the Duke Center for Misophonia and Emotion Regulation.

DISCLOSURE

The authors have nothing to disclose.

REFERENCES

- [1] Swedo SE, Baguley DM, Denys D, et al. Consensus definition of misophonia: a delphi study. *Front Neurosci* 2022;16:841816.
- [2] Brout JJ, Edelstein M, Erfanian M, et al. Investigating misophonia: a review of the empirical literature, clinical implications, and a research agenda. *Front Neurosci* 2018; 12:36.
- [3] Potgieter I, MacDonald C, Partridge L, et al. Misophonia: a scoping review of research. *J Clin Psychol* 2019;75(7): 1203–18.
- [4] Kumar S, Tansley-Hancock O, Sedley W, et al. The brain basis for misophonia. *Curr Biol* 2017;27(4):527–33.
- [5] Edelstein M, Brang D, Rouw R, et al. Misophonia: physiological investigations and case descriptions. *Front Hum Neurosci* 2013;7:296.
- [6] Remmert N, Schmidt KMB, Mussel P, et al. The Berlin Misophonia Questionnaire (BMQ): development and validation of a symptom-oriented diagnostic instrument for the measurement of misophonia. *PsyArXiv* 2021. <https://doi.org/10.31234/osf.io/mujya>.
- [7] Rosenthal MZ, Anand D, Cassiello-Robbins C, et al. Development and initial validation of the duke misophonia questionnaire. *Front Psychol* 2021;12:709928.
- [8] Rosenthal MZ, McMahon K, Greenleaf AS, et al. Phenotyping misophonia: psychiatric disorders and medical health correlates. *Front Psychol* 2022;13:941898.
- [9] Jastreboff MM, Jastreboff PJ. Components of decreased sound tolerance: hyperacusis, misophonia, phonophobia. *ITHS News Letter* 2001;2(5–7):1–5.
- [10] Schröder A, Vulink N, Denys D. Misophonia: diagnostic criteria for a new psychiatric disorder. *PLoS One* 2013; 8(1):e54706.
- [11] Cassiello-Robbins C, Anand D, McMahon K, et al. A Preliminary investigation of the association between misophonia and symptoms of psychopathology and personality disorders. *Front Psychol* 2021;11:519681.
- [12] Cusack SE, Cash TV, Vrana SR. An examination of the relationship between misophonia, anxiety sensitivity, and obsessive-compulsive symptoms. *J Obsessive Compuls Relat Disord* 2018;18:41–8.
- [13] Guetta RE, Cassiello-Robbins C, Trumbull J, et al. Examining emotional functioning in misophonia: the role of affective instability and difficulties with emotion regulation. *PLoS One* 2022;17(2):e0263230.
- [14] Jager I, de Koning P, Bost T, et al. Misophonia: phenomenology, comorbidity and demographics in a large sample. *PLoS One* 2020;15(4):e0231390.
- [15] McKay D, Kim SK, Mancusi L, et al. Profile Analysis of psychological symptoms associated with misophonia: a community sample. *Behav Ther* 2018;49(2):286–94.
- [16] Quek TC, Ho CS, Choo CC, et al. Misophonia in singaporean psychiatric patients: a cross-sectional study. *Int J Environ Res Public Health* 2018;15(7):1410.
- [17] Rouw R, Erfanian M. A large-scale study of misophonia. *J Clin Psychol* 2018;74(3):453–79.

- [18] Siepsiak M, Sobczak AM, Bohaterewicz B, et al. Prevalence of misophonia and correlates of its symptoms among inpatients with depression. *Int J Environ Res Public Health* 2020;17(15):5464.
- [19] Wu MS, Lewin AB, Murphy TK, et al. Misophonia: incidence, phenomenology, and clinical correlates in an undergraduate student sample. *J Clin Psychol* 2014;70(10):994–1007.
- [20] Kılıç C, Öz G, Avanoğlu KB, et al. The prevalence and characteristics of misophonia in Ankara, Turkey: population-based study. *BJPsych Open* 2021;7(5):e144.
- [21] Sheehan DV, Lecrubier Y, Sheehan KH, et al. The mini-international neuropsychiatric interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry* 1998;59(Suppl 20):22–33.
- [22] Jastreboff PJ, Jastreboff MM. Treatments for decreased sound tolerance (hyperacusis and misophonia). *Semin Hear* 2014;35(2):105–20.
- [23] Altınöz AE, Ünal NE, Altınöz ŞT. The effectiveness of cognitive behavioral psychotherapy in misophonia: a case report. *Turk J Clin Psychiatry* 2018;21(4):414–7.
- [24] Bernstein RE, Angell KL, Dehle CM. A brief course of cognitive behavioural therapy for the treatment of misophonia: A case example. *Cogn Behav Ther* 2013;6:e10.
- [25] Cowan EN, Marks DR, Pinto A. Misophonia: a psychological model and proposed treatment. *J Obsessive Compuls Relat Disord* 2022;32:100691.
- [26] Dover N, McGuire JF. Family-based cognitive behavioral therapy for youth with Misophonia: a case report. *Cogn Behav Pract* 2021. <https://doi.org/10.1016/j.cbpra.2021.05.005>.
- [27] Dozier TH. Counterconditioning treatment for Misophonia. *Clin Case Stud* 2015;14(5):374–87.
- [28] Dozier TH. Treating the initial physical reflex of misophonia with the neural repatterning technique: a counterconditioning procedure. *Psychol Thought* 2015;8(2):189–210.
- [29] Kamody RC, Del Conte GC. Using dialectical behavior therapy to treat misophonia in adolescence. *Prim Care Companion CNS Disord* 2017;19(5). <https://doi.org/10.4088/PCC.17102105>.
- [30] Lewin AB, Dickinson S, Kudryk K, et al. Transdiagnostic cognitive behavioral therapy for misophonia in youth: methods for a clinical trial and four pilot cases. *J Affect Disord* 2021;291:400–8.
- [31] McGuire JF, Wu MS, Storch EA. Cognitive-behavioral therapy for 2 youths with misophonia. *J Clin Psychiatry* 2015;76(5):573–4.
- [32] Muller D, Khemlani-Patel S, Neziroglu F. Cognitive-behavioral therapy for an adolescent female presenting with misophonia: a case example. *Clin Case Stud* 2018;17(4):249–58.
- [33] Petersen JM, Twohig MP. Acceptance and commitment therapy for a child with misophonia: a case study. *Clin Case Stud* 2022. <https://doi.org/10.1177/15346501221126115346501221126136>.
- [34] Reid AM, Guzick AG, Gernand A, et al. Intensive cognitive-behavioral therapy for comorbid misophonic and obsessive-compulsive symptoms: a systematic case study. *J Obsessive Compuls Relat Disord* 2016;10:1–9.
- [35] Schneider RL, Arch JJ. Case study: a novel application of mindfulness-and acceptance-based components to treat misophonia. *J Contextual Behav Sci* 2017;6(2):221–5.
- [36] Frank B, McKay D. The suitability of an inhibitory learning approach in exposure when habituation fails: a clinical application to misophonia. *Cogn Behav Pract* 2019;26(1):130–42.
- [37] Jager I, Vulink N, de Roos C, et al. EMDR therapy for misophonia: a pilot study of case series. *Eur J Psychotraumatol* 2021;12(1):1968613.
- [38] Schröder AE, Vulink NC, van Loon AJ, et al. Cognitive behavioral therapy is effective in misophonia: an open trial. *J Affect Disord* 2017;217:289–94.
- [39] Jager IJ, Vulink NC, Bergfeld IO, et al. Cognitive behavioral therapy for misophonia: a randomized clinical trial. *Depress Anxiety* 2020;38(7):708–18.
- [40] Barlow DH, Sauer-Zavala S, Carl JR, et al. The nature, diagnosis, and treatment of neuroticism back to the future. *Clin Psychol Sci* 2014;2(3):344–65.
- [41] Barlow DH, Farchione TJ, Bullis JR, et al. The unified protocol for transdiagnostic treatment of emotional disorders compared with diagnosis-specific protocols for anxiety disorders: a randomized clinical trial. *JAMA Psychiatr* 2017;74(9):875–84.
- [42] Dozier TH, Morrison KL. Phenomenology of misophonia: initial physical and emotional responses. *Am J Psychol* 2017;130(4):431–8.
- [43] Steele SJ, Farchione TJ, Cassiello-Robbins C, et al. Efficacy of the Unified Protocol for transdiagnostic treatment of comorbid psychopathology accompanying emotional disorders compared to treatments targeting single disorders. *J Psychiatr Res* 2018;104:211–6.
- [44] Sakiris N, Berle D. A systematic review and meta-analysis of the Unified Protocol as a transdiagnostic emotion regulation based intervention. *Clin Psychol Rev* 2019;72:101751.
- [45] Cassiello-Robbins C, Sauer-Zavala S, Brody LR, et al. Exploring the effects of the mindfulness and countering emotional behaviors modules from the Unified Protocol on dysregulated anger in the context of emotional disorders. *Behav Ther* 2020;51(6):933–45.
- [46] Tonarely-Busto NA, Phillips DA, Saez-Clarke E, et al. Applying the unified protocol for transdiagnostic treatment of emotional disorders in children and adolescents to misophonia: a case example. *Evid Based Pract Child Adolesc Ment Health* 2022;1–15. <https://doi.org/10.1080/23794925.2022.2025631>.
- [47] McMahon K, Cassiello-Robbins C, Rosenthal MZ. Exploring the Acceptability and Efficacy of a Transdiagnostic Treatment for Misophonia. *Misophonia Research Fund Annual Report* 2021;1–3.
- [48] S.C. Hayes, S.G. Hofmann (Eds.), *Process-based CBT: the science and core clinical competencies of cognitive behavioral therapy*, New Harbinger Publications, Inc, Oakland, CA, 2018.
- [49] Jakubovski E, Müller A, Kley H, et al. Prevalence and clinical correlates of misophonia symptoms in the general population of Germany. *Front Psychiatry* 2022;13:012424.